

## **Lorandite from Allchar, a long-term double detector for pp-neutrinos and fast myons**

### **Lorandite from Allchar, a long-term double detector for pp-neutrinos and fast myons**

**Pavicevic M. K.<sup>1</sup>, Amthauer G.<sup>1</sup>, Anicin I.<sup>2</sup>, Bosch F.<sup>3</sup>, Boev B.<sup>4</sup> and Pejovic V.<sup>2</sup>**

<sup>1</sup> University of Salzburg, faculty of material engineering & physics, A-5020 Salzburg

<sup>2</sup> University of Belgrade, faculty of physics, RS-11000 Belgrade

<sup>3</sup> GSI – Gesellschaft für Schwer Ionenforschung mbH, D-64220 Darmstadt

<sup>4</sup> University of Stip, faculty of mining and geology, MK-92000 Stip

The Sb-As-Tl-Au deposit at Allchar, providing one of the world-wide largest concentrations of thallium and also numerous other Tl-bearing minerals, may open an outstanding scientific perspective. Allchar belongs to the Serbian – Macedonian metallogenic province located near the border between Macedonia and Greece. By a long-lasting international research it has been shown that Lorandite from Allchar can serve, together with cogenetic monitor minerals, as a geochemical detector for both, the flux of solar pp-neutrinos, averaged over the geological age of Lorandite, and for the average flux of fast cosmic myons within the same period of time.

By the capture of (mainly) solar pp-neutrinos with an unprecedented low threshold of only 52 keV for them, <sup>205</sup>Tl is transformed to <sup>205</sup>Pb. Our investigations show that about 22 atoms <sup>205</sup>Pb in 1g Lorandite for the geological time range of 4.2 Ma and a palaeozoic depth of 570 m should be expected, by supposing the present solar neutrino luminosity and by taking into account its reduction due to neutrino flavour oscillations. However, an additional production of <sup>205</sup>Pb occurs via the interaction of cosmic radiation (stopped and fast myons) with decay products of <sup>238</sup>U and <sup>232</sup>Th. The amount of this "underground" of <sup>205</sup>Pb depends on the palaeozoic depth (actual depth plus eroded sheets) and the geological age of the Tl-mineralization. Calculations based on known nuclear cross-sections show that fast myons generate by far the largest part of "underground" <sup>205</sup>Pb atoms. The present state of research predicts, for palaeozoic depths of 350 m and 570 m, respectively, total numbers of 106 and 48 <sup>205</sup>Pb atoms per gram of Lorandite for 4.2 Ma, whereby the contributions from fast myons add up to 84 and 26 atoms of <sup>205</sup>Pb, respectively. For the detection of the few <sup>205</sup>Pb atoms we will apply SMS (Schotty Mass Spectrometry) at the ion storage ring of GSI (Gesellschaft für Schwerionenforschung, Darmstadt), where single fully-stripped <sup>205</sup>Pb ions can be detected.

References:

1. Proceedings of the International Symposium on Solar Neutrino Detection with <sup>205</sup>Tl, Dubrovnik, Yugoslavia 1990, ed. By M. K. Pavicevic and G. Amthauer, Beih. N. Jb. Miner. Abh. 2/3, 125 (1994)

Abs. No. **560**

Meeting: **DMG 2008**

submitted by: **Pavicevic, Miodrag**

email: **miodrag.pavicevic@sbg.ac.at**

date: **0000-00-00**

Req. presentation: **Poster**

Req. session: **S02**